

The Effects of Blood Glucose Control in the Operative Spine Patient: A Systematic Review

Richard Bidwell, Michael Spitnale, Rodrigo Encinas, Yianni Bakaes, Justin Kung and Gregory Grabowski

Int J Spine Surg published online 12 October 2023
<https://www.ijssurgery.com/content/early/2023/10/09/8547>

This information is current as of February 2, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

The Effects of Blood Glucose Control in the Operative Spine Patient: A Systematic Review

RICHARD BIDWELL, MD¹; MICHAEL SPITNALE, MD¹; RODRIGO ENCINAS, MD²; YIANNI BAKAES, BS²; JUSTIN KUNG, MD¹; AND GREGORY GRABOWSKI, MD¹

¹Department of Orthopaedic Surgery, Prisma Health Midlands, Columbia, SC, USA; ²Department of Orthopaedic Surgery, University of South Carolina School of Medicine Columbia, Columbia, SC, USA

ABSTRACT

Objective: To our knowledge, this is the first systematic review to evaluate the available literature on the effects of perioperative serum glucose (SG) on outcomes for patients undergoing spine surgery. This review will add insight into how the perioperative management of SG affects the outcomes of patients undergoing spine surgery.

Methods: Three databases were used in this review including Embase, PubMed, and Cochrane Library. The searches were from 2012 to 2022 and included the terms “spine surgery” and “glucose level” to identify studies that demonstrated a correlation between glucose level and postoperative outcomes. Pediatric studies, those that did not specify spine surgical outcomes related to glucose levels, and non-English studies were excluded. The methodological items for nonrandomized studies score was used to assess risk of bias in the included studies.

Results: This review included a total of 9 cohort studies, both prospective and retrospective, encompassing a total of 431,156 subjects. Seven of the 9 studies reported an increased overall complication rate among patients with diabetes or with higher SG levels, and 4 studies demonstrated an increased infection rate among this population. Two studies reported an association between decreased SG levels and improved neurological recovery when a deficit was present preoperatively, and 1 of the studies found that this association was statistically significant.

Limitations: Limitations of this review include lack of standardization regarding type of surgery, location of the spine, and level of evidence.

Conclusion: Most of the current literature suggests that elevated SG levels in patients undergoing spine surgery likely leads to higher complication rates and may lead to increased infection rates, and this review reinforced the current evidence. Additionally, perioperative SG levels may be associated with the extent of neurological recovery after surgery, but further investigation may be warranted.

Clinical Relevance: This review adds to the current body of evidence regarding perioperative SG levels and its association with complications.

Level of Evidence: 2.

Complications

Keywords: spine surgery, glucose control, outcomes, diabetes

INTRODUCTION

Spine surgery is one of the most common surgical procedures performed in the United States.¹ Traumatic injuries, degenerative processes, and other medical conditions lead to numerous cervical, thoracic, and lumbar procedures each year. Pathologies treated range from disc herniations to major deformity correction and stabilization of traumatic injuries. Given this, there has been an observed increase in the number of spine surgeries worldwide. Kobayashi et al reported data from 10 facilities in Japan and showed a 2.4 times increase in spine cases over the past 15 years.² It is worth mentioning that Japan has a predominant aging population. Norway is another country with an abundant aging population, and Grotle et al reported an increase of 54% in spine surgeries in Norway over a 15-year period.³

The population in the United States is aging as well, and there has been a recent increase in the need for operative intervention for spine pathology.⁴ A recent analysis by Martin et al reported a 62.3% increase in lumbar fusion surgery in the United States over an 11-year period, with the highest number of procedures performed in people aged 65 years or older.⁵ Additionally, Deng et al reported an increase greater than 250% for lumbar fusion surgery among patients with degenerative disease over a 12-year period in the United States.⁶

Given the notable increase in operative spine patients, it may be important to consider factors that can improve surgical outcomes. Diabetes is strongly correlated to obesity, and since obesity and diabetes are increasing among the population at an alarming rate, there may be an increased need for spine surgeries among this population.^{7,8}

Diabetes and uncontrolled glucose levels have been previously reported to negatively affect outcomes in orthopedic surgery.⁹ Browne et al reviewed a retrospective cohort and reported that patients with diabetes had a statistically significant higher incidence of mortality and infection after lumbar spine surgery.¹⁰ Another retrospective study by Chen et al reported that the strongest risk factor for developing infection after lumbar spine surgery was diabetes.¹¹

On the other hand, some literature showed no difference among diabetic patient outcomes after undergoing spine surgery. Of note, these studies analyzed non-insulin-dependent diabetics. Cho et al reported no significant difference in complications after 2 years among non-insulin-dependent diabetic patients and patients without diabetes undergoing lumbar spine surgery.¹² However, this study had a small sample size of only 42 patients, which may limit the ability of their results to be generalizable to all surgical patients. Additionally, Dokai et al reported no significantly different major complications among diabetic vs nondiabetic patients undergoing cervical spine surgery for cervical myelopathy.¹³ The only significant differences between the diabetic patients vs nondiabetic patients in this study were worse recovery of motor and sensory function in the lower extremities in diabetic patients and a negative correlation between recovery rate and preoperative HbA1c. However, the results of this study should be taken cautiously considering its limitations of a small sample size of only 78 patients and its retrospective nature.

In addition to elective spine surgeries, prior literature suggests that abnormal blood glucose levels may significantly worsen outcomes in the acute setting of spinal cord injuries. Furlan et al retrospectively evaluated a cohort of hyperacute traumatic spinal cord injuries and abnormalities in blood glucose among 499 patients.¹⁴ Blood glucose levels >140 mg/dL after 7 days of injury were associated with a statistically significant increase in mortality and a decrease in the Functional Independence Measure score within 1 year.

To our knowledge, this is the first systematic review to evaluate the available literature on the effects of diabetes mellitus and abnormalities in glucose levels on outcomes for patients undergoing spine surgery. This systematic review will add significant evidence to the current literature and improve the preoperative and postoperative management of patients undergoing spine surgery.

MATERIALS AND METHODS

This systematic review was designed and conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analysis protocol and the Cochrane Handbook guidelines.^{15,16} There was no required

institutional board review approval for this systematic review.

Search Strategy

Three databases were used in this systematic review: Embase, PubMed, and Cochrane Library. The searches were from 2012 to 2022 and included the following terms: “spine surgery” and “glucose level”. Initial screening of titles, abstracts, and full-text studies was performed.

Inclusion and Exclusion Criteria

The inclusion criteria were any studies published between 2012 and 2022 that included patients undergoing spine surgery and had a correlation between glucose level and postsurgical outcomes. Pediatric studies, those that did not specify spine surgical outcomes related to glucose levels, and non-English studies were excluded.

Data Collection and Abstraction

The investigators independently evaluated titles, abstracts, and full text from the studies found in the search. The senior author reviewed the available studies and resolved any discussion from the search, inclusion, and exclusion criteria for the studies. The following data were extracted: authors, the title of studies, journal of publication, year of publication, country, level of evidence, number of subjects, number of procedures performed, mean age, number of women and men, visual analog scale (VAS) scores, final follow-up time, control group, surgical procedures, outcomes, and complications.

Statistical Analysis Assessment for Risk of Bias

The methodological items for nonrandomized studies score was used to evaluate the quality of the studies (Table 1).¹⁷ All 9 studies adequately reported a clear aim, a prospective collection of data, appropriate end points of the studies, inclusion of consecutive patients, follow-up periods, an adequate statistical analysis, prospective calculations of the study size, and loss of follow-up less than 5% except for the study by Tanashima et al.¹⁸ The study by Peng et al was the only one to earn a score of 2 for an unbiased assessment of the study endpoint; the rest of the studies did not report it.¹⁹ Four studies reported having control groups.^{18,20–22} The average score of the studies included in this systematic review was 17.7 with a range from 14 to 22. These scores represent high-quality studies.

Table 1. MINORS criteria scores by study.

Criteria	Guzman et al ²³	Peng et al ¹⁹	Tanishima et al ¹⁸	Hwang et al ²⁴	Worley et al ²⁵	Langlois et al ²⁶	Khan et al ²⁰	Moazzeni et al ²²	Machino et al ²¹
A clearly stated aim	2	2	2	2	2	2	2	2	2
Inclusion of consecutive patients	2	2	2	2	2	2	2	2	2
Prospective collection of data	2	2	2	2	2	2	2	2	2
Endpoints appropriate to the aim of the study	2	2	2	2	2	2	2	2	2
Unbiased assessment of the study endpoint	0	2	0	0	0	0	0	0	0
Follow-up period appropriate to the aim of the study	0	0	2	2	0	2	2	2	2
Loss to follow-up less than 5%	2	2	1	2	2	2	2	2	2
Prospective calculation of the study size	2	2	2	2	2	2	2	2	2
An adequate control group	0	0	1	0	0	0	1	2	2
Contemporary groups	0	0	2	0	0	0	1	2	2
Baseline equivalence of groups	0	0	2	0	0	0	1	2	2
Adequate statistical analyses	2	2	2	2	2	2	2	2	2
Total score	14	16	20	16	14	16	19	22	22

Abbreviation: MINORS, methodological items for nonrandomized studies.

Note: MINORS Score Criteria: scored 0 (not reported), 1 (reported but inadequate), or 2 (reported and adequate).²³

RESULTS

Characteristics of the Systematic Review Search and Studies

A total of 1083 results were found after the primary databases were searched. PubMed included 178 results, Embase included 850 results, and Cochrane Library included 55 studies. A total of 630 records were removed after adjusting for the selected date range. From the remaining 453 studies, 19 studies were excluded due to non-English language (2) and lack of full-text availability (17). A total of 434 full-text articles were assessed for eligibility, and 425 were excluded due to not meeting the criteria for inclusion or being repeated from other database results. Thus, a total of 9 studies were included in this systematic review, which included a total of 431,156 patients (Figure).

The 9 studies we analyzed included data from 6 countries, with studies and subjects from the United States being most represented. Of the studies that specified age, the average age ranged from 56.1 to 69.5 years. The percentage of women in these studies ranged from 43% to 65.2%. Five of these studies included lumbar spine surgery, 3 of them included cervical spine surgery, and 1 study included cervical, thoracic, or lumbar spine surgery. Additionally, 5 of these studies were retrospective and 4 were prospective (Table 2).^{18–26}

Postsurgical Outcomes by Study

The retrospective database analysis by Guzman et al came from the Nationwide Inpatient Sample database and included 3 types of procedures: lumbar decompression without fusion, lumbar fusion, and lumbar

fusion revision.²³ Among all subjects, the group with uncontrolled/controlled diabetes carried a higher risk of having other comorbidities such as congestive heart failure, chronic pulmonary disease, liver disease, neurological disorder, renal disease, obesity, and depression compared with subjects without diabetes. The separation criteria for inclusion into either group was based on




Figure. Flow diagram for search strategy and selection criteria.

Table 2. Demographics and characteristics of the studies.

Author	Journal	Year	Country	Level of Evidence	Subjects	Mean Age, y	Type of Procedure	No. of Procedures	% Women	Follow-Up
Guzman et al ²³	Spine	2014	USA	III: Retrospective database analysis	423,050	61.8	Lumbar spine surgery	2,568,994	51.8%	NR
Peng et al ¹⁹	Medicine	2019	China	III: Retrospective single-center analysis	523	NR ^a	Lumbar spine surgery	523	43.6%	NR
Tanishima et al ¹⁸	Asian Spine Journal	2019	Japan	II: Multicenter prospective study	61	69.5	Cervical spine surgery	61	43%	12 mo
Hwang et al ²⁴	Korean Journal of Neurotrauma	2019	Korea	III: Retrospective review	92	68.4	Lumbar spine surgery	92	58.7%	23.4 mo
Worley et al ²⁵	International Journal of Spine Surgery	2017	USA	III: Retrospective review	5904	61	Cervical spine surgery	5904	42.6%	NR
Langlois et al ²⁶	European Spine Journal	2014	France	II: Prospective cohort study	75	56.1	Cervical, lumbar, and thoracic spine surgery	75	62.7%	3 mo
Khan et al ²⁰	Global Spine Journal	2019	USA	III: Retrospective cohort study	850	58.4	Lumbar spine surgery	850	53.3%	22.99 mo
Moazzeni et al ²²	International Journal of Spine Surgery	2018	Iran	II: Prospective study	96	57.8	Lumbar spine surgery	96	65.2%	12 mo
Machino et al ²¹	Spine	2017	Japan	II: Prospective cohort study	505	66.6	Cervical spine surgery	505	39.2%	12 mo

Abbreviation: NR, not reported.

^aThree hundred thirty patients were older than 60 y.

ICD-9-CM codes. There was a statistically significantly increased risk of postoperative infection ($P < 0.0001$), inpatient mortality ($P = 0.0001$), postsurgical bleeding, and revision rate among patients with diabetes vs patients without diabetes. Among those with diabetes, patients with uncontrolled diabetes had an increased risk of postoperative infection and inpatient mortality compared with those with controlled diabetes. Length of stay ($P < 0.0001$) was also significantly higher in patients with diabetes (regardless of being controlled) vs patients without diabetes.

The retrospective, single-center analysis by Peng et al reported outcomes of surgical intervention for lumbar stenosis, lumbar disc herniation, lumbar spondylolisthesis, and lumbar vertebral fractures.¹⁹ Their patient sample only included patients with diabetes mellitus. Only 37 patients (7.1%) developed a postoperative infection. Their results from a univariate analysis showed the following statistically significant χ^2 test for developing infection: a 10.67 times increased risk ($P = 0.01$) for patients with preoperative HbA1c $> 7.5\%$ compared with those with $< 7.5\%$, and an 18.57 times increased risk ($P < 0.001$) at 3 days postoperatively with a fasting blood glucose level > 144 mg/dL (3 days average of all glucose measurements) compared with those with levels < 144 mg/dL. There was also an observed 23.79 times increased risk ($P < 0.001$) at 3 days postoperatively with an average postprandial blood glucose > 180 mg/dL compared with those patients with a 3-day postprandial glucose average < 180 mg/dL. Their results from the multivariate analysis showed a statistically significant odds ratio of 4.63 for developing infection with a preoperative HbA1c

$> 7.5\%$, an odds ratio of 5.71 for 3-day postoperative fasting blood glucose > 144 mg/dL, and an odds ratio of 7.74 for 3-day postoperative average postprandial blood glucose > 180 mg/dL.

The multicenter prospective study by Tanishima et al analyzed laminoplasty and anterior decompression and fusion for cervical spondylotic myelopathy and ossification of the posterior longitudinal ligament.¹⁸ They reported no significant differences in postoperative improvement among patients with diabetes vs without diabetes. At 1-year follow-up, the diabetic patients with good control of HbA1c (undefined in the study methods) had a statistically significant ($P < 0.01$) improvement in upper and lower limb function based on the previously validated Japanese Orthopedic Association Myelopathy Evaluation Questionnaire. However, there was no statistically significant difference when comparing the 2 groups. It is worth mentioning that this study had the lowest number of subjects (61) in this review, consisting of 0.01% of subjects in our study. Another limitation of this study was the high rate of loss to follow-up, with only 61 patients out of the 140 (43.6%) available for 1-year follow-up.

The retrospective review by Hwang et al analyzed single-level posterolateral and posterior interbody fusion surgery.²⁴ They evaluated the relationship between HbA1c and risk of infection and reported that preoperative HbA1c was statistically significantly higher in patients who developed a postoperative infection vs patients who did not develop an infection (HbA1c of 6.78% vs 5.99%, respectively, $P = 0.003$). A surgical site infection was defined as having a superficial infection within 30 days of surgery or as having a deep infection within 1 year of surgery. The

incidence of infection was 26.1%, with 3.3% in deep infections and 22.8% in superficial infections. Six infections resulted from the posterolateral approach, and 18 infections resulted from the posterior interbody fusion. The univariate and multivariate logistic analyses results showed that an HbA1c greater than 6.9% was an independent risk factor for infection with an odds ratio of 4.5 ($P = 0.008$). Additionally, longer periods of hospitalization were statistically significant, with 45.8 days in the hospital for patients who had a postoperative infection vs 19.3 days for patients who did not ($P = 0.003$).

The retrospective review by Worley et al analyzed the surgical treatment for patients with cervical spondylotic myelopathy.²⁵ The patient samples included nondiabetic, insulin-independent diabetic, and insulin-dependent diabetic patients. Patients with diabetes were more likely to have cardiac and pulmonary comorbidities and higher body mass index values. A statistically significant increase in pneumonia, unplanned intubation, ventilator for more than 48 hours, bleeding transfusion, and urinary tract infections among patients with diabetes was reported. Longer hospital stays were found among patients with diabetes ($P < 0.001$). The only statistically significant difference found among diabetic patients was an increased incidence of wound complications ($P < 0.001$) for insulin-dependent patients vs for noninsulin-dependent patients (2.1% vs 0.3%, respectively, $P = 0.045$).

The prospective cohort study by Langlois et al analyzed surgeries of the cervical, thoracic, and lumbar levels of the spine.²⁶ They evaluated levels of glucose among nondiabetic patients and found a 29.3% incidence of postoperative complications within 3 months. There were statistically significantly higher glucose levels during the first 3 days of surgery when compared with preoperative values. On postoperative day 2, the average blood glucose level for patients who needed surgical revision was 140.4 ± 16.2 mg/dL, which was significantly higher than that of patients who did not need a revision (blood glucose level: 120.6 ± 21.6 mg/dL; $P = 0.04$). Factors that may have been associated with this increase in glucose included age, smoking status, instrumentation, operative time, blood loss, and need for a transfusion.

The retrospective cohort study by Khan et al analyzed open posterior lumbar fusion for degenerative spondylolisthesis.²⁰ They compared outcomes between diabetic vs nondiabetic patients. The only reported statistically significant difference among these groups was higher discharge rates to facilities for diabetic patients ($P = 0.018$). Radiological measures showed a statistically significantly higher pelvic incidence-lumbar lordosis difference for patients with diabetes. There were no statistically significant differences

among the groups regarding postoperative complications, VAS scores, reoperation, or length of hospital stay.

The prospective study by Moazzeni et al analyzed lumbar fusion for patients with canal stenosis, disc herniation, and degenerative spondylolisthesis.²² They evaluated surgical outcome differences among diabetic vs nondiabetic patients. Fusion status was assessed by lumbar computed tomography scan and anteroposterior and lateral x-rays images 1-year postoperatively. They reported a statistically significant increase in fusion rate in nondiabetic patients at 78% vs in diabetic patients at 53% after 1-year ($P = 0.02$). Additionally, at the last follow-up, statistical significance was observed to be higher in diabetic patients for both Oswestry Disability Index (ODI) scores (average score in diabetic patients was 52 vs average score in nondiabetic patients was 41.3 [$P < 0.001$]), and for postoperative bleeding ($P = 0.006$). VAS scores were significantly higher preoperatively in diabetic patients ($P = 0.07$); however, the postoperative scores were not. Among diabetic patients, insulin-dependent diabetics had statistically significantly higher HbA1c, preoperative ODI, and 6-month postoperative ODI.

The prospective cohort study by Machino et al analyzed cervical double door laminoplasty in patients with cervical spondylotic myelopathy.²¹ They evaluated both diabetic and nondiabetic patients and compared their outcomes after 1 year postoperatively. The diabetic patients were classified as having a previous diagnosis or having fasting blood glucose >126 mg/dL. All patients were confirmed with an HbA1c test before meeting criteria for the diabetic group. It was reported that the recovery rates of lower extremity motor function ($P = 0.047$), upper extremity sensory function ($P = 0.006$), and urinary bladder function ($P = 0.035$) were significantly lower in diabetic patients. It was also reported that all patients had well-controlled glucose levels during the perioperative period. The study did not report differences among patients with a history of diabetes vs those who were recently diagnosed and did not report whether patients' glucose levels were monitored during the 1-year follow-up period. There was no significant difference among recovery rates for upper extremity motor function, lower extremity sensory function, and trunk sensory function. The preoperative and postoperative results of the 10-second step test were significantly lower in the diabetic group ($P < 0.001$). Machino et al did not report on the rate of infection. There was no significant difference between the 2 patient groups regarding operative time, blood loss, postoperative cervical alignment, or range of motion.

The overall number of postoperative complications per study is shown in Table 3. All comparisons were based on diabetic subjects vs nondiabetic subjects, or among patients

Table 3. Complications per study.

Complications	Guzman et al ²³	Peng et al ¹⁹	Tanishima et al ¹⁸	Hwang et al ²⁴	Worley et al ²⁵	Langlois et al ²⁶	Khan et al ²⁰	Moazzeni et al ²²	Machino et al ²¹
Infection	Increased	Increased	NR	Increased	Increased	NR	ND	ND	NR
Mortality	Increased	NR	NR	NR	NR	NR	NR	NR	NR
JOAMRQ	NR	NR	ND	NR	NR	NR	NR	NR	NR
Longer hospitalization	Increased	NR	NR	NR	Increased	NR	ND	ND	NR
PI-LL	NR	NR	NR	NR	NR	NR	Increased	NR	NR
Fusion rate	NR	NR	NR	NR	NR	NR	NR	Decreased	NR
Postoperative bleeding	Increased	NR	NR	NR	Increased	NR	NR	Increased	ND
Oswestry Disability Index	NR	NR	NR	NR	NR	NR	ND	Increased	NR
Motor function recovery	NR	NR	ND	NR	NR	NR	NR	NR	Decreased
Sensory function recovery	NR	NR	ND	NR	NR	NR	NR	NR	Decreased
Bladder function recovery	NR	NR	ND	NR	NR	NR	NR	NR	Decreased
Surgical revision	Increased	NR	NR	NR	NR	Increased	ND	NR	NR

Abbreviations: JOAMRQ, Japanese Orthopedic Association Myelopathy Evaluation Questionnaire; ND, no difference; NR, not reported; PI-LL, pelvic incidence-lumbar lordosis difference.

with higher levels of glucose vs those with normal levels of glucose. Only statistically significant differences were reported. The studies by Machino et al, Moazzeni et al, and Tanishima et al did not report specific differences on the values for glucose levels or HbA1c.^{18,21,22} Guzman et al reported on controlled diabetic patients, uncontrolled diabetic patients, and nondiabetic patients; however, this study did not include specific blood glucose or HbA1c values.²³ Worley et al reported on diabetic patients stratified based on whether their diabetes was insulin or non-insulin dependent, but no specific glucose values were noted in this study.²⁵ Khan et al reported an average HbA1c of 7.2% (range 5.1–10.7) for diabetic patients in their study.²⁰ Langlois et al reported on nondiabetic patients and found that, on postoperative day 2, the average glucose level for patients who needed surgical revision was 140.4 ± 16.2 mg/dL, which was significantly higher than that of patients who did not need a revision, with an average blood glucose level of 120.6 ± 21.6 mg/dL ($P = 0.04$).²⁶ Peng et al reported that values of postprandial glucose >180 mg/dL, postoperative fasting glucose >144 mg/dL, and/or HbA1c >7.5 were associated with a higher rate of infection.¹⁹ Hwang et al reported that preoperative HbA1c was higher for patients who developed a postoperative infection (6.78%) vs those who did not develop an infection (5.99%; $P = 0.003$). Additionally, they found that an HbA1c $>6.9\%$ was associated with a significant increase in surgical site infection ($P = 0.001$).¹²

DISCUSSION

The goal of this systematic review was to evaluate and analyze the current available literature on the outcomes of diabetes and abnormalities in blood glucose levels for patients undergoing spine surgery. During our search, we were unable to find any other meta-analysis or systematic review on this specific topic. Spine surgery is one of

the most frequently performed orthopedic surgeries in the United States. The literature also demonstrates a considerable increase in the number of patients worldwide undergoing spine surgery in recent years, particularly in countries with a large aging population.^{2,3,5,6}

This systematic review included 9 studies that consisted of both prospective and retrospective studies. There were no randomized controlled trials included. Among the included studies were 4 studies that reported an increase in the rate of infection among patients with diabetes or higher levels of glucose.^{19,23–25} There were 2 studies that reported no significant difference in the rate of infection among patients with diabetes vs patients without diabetes.^{20,22} Three studies did not report information regarding infection.^{18,21,26} The 4 studies that included a statistically significantly higher rate of infection included a total of 429,569 subjects, which represented 99.6% of the patients included in this systematic review.^{19,23–25} There was only 1 study that reported a mortality risk, which was statistically significantly higher in patients with diabetes compared with that of patients without diabetes.²³ This study was a database analysis that included 98.1% of patients in this review.

Overall, 7 of the 9 studies included in this review reported a statistically significantly higher complication rate among patients with diabetes or abnormalities in blood glucose levels compared with that of patients with normal blood glucose levels. Tanishima et al did not report an increase in postoperative complications among diabetic patients, but they included only 61 subjects and had only 1-year postoperative follow-up data for 43.6% of patients analyzed.¹⁸ Machino et al reported on operative time and blood loss values but did not investigate complications such as infection.²¹

An interesting finding from this review came from the study by Machino et al that analyzed cervical spondylotic myelopathy patients, demonstrating a delayed neurological

recovery rate in diabetic patients or in patients who had uncontrolled postoperative blood glucose.²¹ This phenomenon was not found to be well studied during our literature review. It appears that there is currently very limited data regarding outcomes of neurological recovery in the surgically managed spinal trauma patient population or in patients with cervical spondylotic myelopathy. Tanishima et al investigated neurological recovery in patients with the same diagnosis and showed a trend toward improved recovery with better serum glucose (SG) control, but they did not find a statistically significant difference.¹⁸ From the data found in this review regarding higher rates of wound infections and medical complications in diabetic patients, one may expect similarly worsened outcomes for neurologic recovery, but this topic may require further exploration. It may also be worth researching whether more rigorous blood glucose control plays a role in neurological recovery after a traumatic spinal cord injury that is managed operatively. Given that patients who sustain traumatic spine injuries that meet indications for operative intervention are often polytrauma patients, it would be advantageous to understand whether we can significantly improve their overall outcomes, recovery, and mortality by tightly controlling blood glucose levels.

One limitation of this review was the lack of reporting on the use of intrawound antibiotic powder that is becoming common practice. How this plays a role in the reduction of infection for patients with better perioperative glucose control also remains to be seen. A recent study by Ushirozako et al included more than 1200 patients and found a decrease in the number of surgical site infections in both superficial and deep tissue in the vancomycin group vs the control group, but this was not statistically significant.²⁷ However, when Ushirozako et al used baseline and surgical data of propensity score–matched patients, they found a statistically significant decrease in the occurrence of infection by half for the vancomycin group ($P = 0.041$). Additionally, a meta-analysis by Tailaiti et al included 27 studies with more than 17,000 subjects and reported that vancomycin significantly reduced the incidence of surgical site infection for those undergoing internal fixation of the spine ($P = 0.01$).²⁸ Although patients with diabetes and abnormalities in blood glucose were included in these studies, there were no specifications made regarding the ability of vancomycin to improve outcomes based on the severity of their disease/ blood glucose levels.

Another limitation of this review is the lack of standardization of the included studies regarding type of surgery, location of the spine, whether trauma patients were included, and level of evidence, as these factors can significantly alter results. Also, there is potential for human error

in the reported SG levels because humans were responsible for performing the test and appropriately documenting this information. Additionally, we were unable to account for the possibility of patients receiving steroids, which are known to impact neurological deficits, pain control, control of dysphagia symptoms, and SG levels, in the perioperative period.

CONCLUSION

This systematic review will add to the current body of evidence regarding perioperative SG levels and their association with postoperative complications and infections. Most of the current literature suggests that elevated SG levels in patients undergoing spine surgery may lead to postoperative infections, and this review reinforced the current evidence. In addition, perioperative SG levels may be associated with the extent of neurological recovery after spine surgery when a deficit is present preoperatively, but further investigation may be warranted.

REFERENCES

1. Fingar KR, Stocks C, Weiss AJ, Steiner CA. Most Frequent Operating Room Procedures Performed in U.S. Hospitals, 2003-2012. Most Frequent Operating Room Procedures Performed in U.S. Hospitals, 2003-2012 #186. Published December 2014. 2003. <https://www.hcup-us.ahrq.gov/reports/statbriefs/sb186-Operating-Room-Procedures-United-States-2012.jsp>. Accessed July 2022.
2. Kobayashi K, Sato K, Kato F, et al. Trends in the numbers of spine surgeries and spine surgeons over the past 15 years. *Nagoya J Med Sci.* 2022;84(1):155–162. doi:10.18999/nagjms.84.1.155
3. Grotle M, Småstuen MC, Fjeld O, et al. Lumbar spine surgery across 15 years: trends, complications and reoperations in a longitudinal observational study from Norway. *BMJ Open.* 2019;9(8):e028743. doi:10.1136/bmjopen-2018-028743
4. Vespa J, Medina L, Armstrong DM. Demographic turning points for the United States... - census.gov. Published March 2018. <https://www.census.gov/content/dam/Census/library/publications/2020/demo/p25-1144.pdf>. Accessed July 22, 2022.
5. Martin BI, Mirza SK, Spina N, Spiker WR, Lawrence B, Brodke DS. Trends in lumbar fusion procedure rates and associated hospital costs for degenerative spinal diseases in the United States. *Spine.* 2019;44(5):369–376. doi:10.1097/BRS.0000000000002822
6. Deng H, Yue JK, Ordaz A, Suen CG, Sing DC. Elective lumbar fusion in the United States: national trends in inpatient complications and cost from 2002-2014. *J Neurosurg Sci.* 2021;65(5):503–512. doi:10.23736/S0390-5616.19.04647-2
7. Agha M, Agha R. The rising prevalence of obesity: part A: impact on public health. *Int J Surg.* 2017;2(7):e17–e17. doi:10.1097/IJ9.0000000000000017
8. Rowley WR, Bezold C, Arikan Y, Byrne E, Krohe S. Diabetes 2030: insights from yesterday, today, and future trends. *Popul Health Manag.* 2017;20(1):6–12. doi:10.1089/pop.2015.0181
9. Wukich DK. Diabetes and its negative impact on outcomes in orthopaedic surgery. *World J Orthop.* 2015;6(3):331–339. doi:10.5312/wjo.v6.i3.331

10. Browne JA, Cook C, Pietrobon R, Bethel MA, Richardson WJ. Diabetes and early postoperative outcomes following lumbar fusion. *Spine (Phila Pa 1976)*. 2007;32(20):2214–2219. doi:10.1097/BRS.0b013e31814b1bc0
11. Chen S, Anderson MV, Cheng WK, Wongworawat MD. Diabetes associated with increased surgical site infections in spinal arthrodesis. *Clin Orthop Relat Res*. 2009;467(7):1670–1673. doi:10.1007/s11999-009-0740-y
12. Cho W, Lenke LG, Bridwell KH, et al. Comparison of spinal deformity surgery in patients with non-insulin-dependent diabetes mellitus (NIDDM) versus controls. *Spine (Phila Pa 1976)*. 2012;37(16):E978–E984. doi:10.1097/BRS.0b013e31824edf42
13. Dokai T, Nagashima H, Nanjo Y, Tanida A, Teshima R. Surgical outcomes and prognostic factors of cervical spondylotic myelopathy in diabetic patients. *Arch Orthop Trauma Surg*. 2012;132(5):577–582. doi:10.1007/s00402-011-1449-4
14. Furlan JC. Effects on outcomes of hyperglycemia in the hyperacute stage after acute traumatic spinal cord injury. *Neurotrauma Rep*. 2021;2(1):14–24. doi:10.1089/neur.2020.0042
15. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71:71. doi:10.1136/bmj.n71
16. Higgins JPT, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions*. 2nd Edition. Chichester (UK): John Wiley & Sons; 2019. <https://onlinelibrary.wiley.com/doi/book/10.1002/9781119536604>. doi:10.1002/9781119536604
17. Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (MINORS): development and validation of a new instrument. *ANZ J Surg*. 2003;73(9):712–716. doi:10.1046/j.1445-2197.2003.02748.x
18. Tanishima S, Mihara T, Tanida A, et al. Influence of diabetes mellitus on surgical outcomes in patients with cervical myelopathy: a prospective, multicenter study. *Asian Spine J*. 2019;13(3):468–477. doi:10.31616/asj.2018.0082
19. Peng W, Liang Y, Lu T, et al. Multivariate analysis of incision infection after posterior lumbar surgery in diabetic patients: a single-center retrospective analysis. *Medicine (Baltimore)*. 2019;98(23):e15935. doi:10.1097/MD.0000000000015935
20. Khan JM, Michalski J, Basques BA, et al. Do clinical outcomes and sagittal parameters differ between diabetics and nondiabetics for degenerative spondylolisthesis undergoing lumbar fusion? *Global Spine J*. 2020;10(3):286–293. doi:10.1177/2192568219850090
21. Machino M, Imagama S, Ando K, et al. Characteristics of residual symptoms after laminoplasty in diabetic patients with cervical spondylotic myelopathy: a prospective cohort study. *Spine (Phila Pa 1976)*. 2017;42(12):E708–E715. doi:10.1097/BRS.0000000000001947
22. Moazzeni K, Kazemi KA, Khanmohammad R, Eslamian M, Rostami M, Faghih-Jouibari M. Comparison of surgical outcome between diabetic versus nondiabetic patients after lumbar fusion. *Int J Spine Surg*. 2018;12(4):528–532. doi:10.14444/5064
23. Guzman JZ, Iatridis JC, Skovrlj B, et al. Outcomes and complications of diabetes mellitus on patients undergoing degenerative lumbar spine surgery. *Spine (Phila Pa 1976)*. 2014;39(19):1596–1604. doi:10.1097/BRS.0000000000000482
24. Hwang JU, Son DW, Kang KT, et al. Importance of hemoglobin A1C levels for the detection of post-surgical infection following single-level lumbar posterior fusion in patients with diabetes. *Korean J Neurotrauma*. 2019;15(2):150–158. doi:10.13004/kjnt.2019.15.e36
25. Worley N, Buza J, Jalai CM, et al. Diabetes as an independent predictor for extended length of hospital stay and increased adverse post-operative events in patients treated surgically for cervical spondylotic myelopathy. *Int J Spine Surg*. 2017;11(2):10. doi:10.14444/4010
26. Langlois J, Bouyer B, Larroque B, Dauzac C, Guigui P. Glycemic instability of non-diabetic patients after spine surgery: a prospective cohort study. *Eur Spine J*. 2014;23(11):2455–2461. doi:10.1007/s00586-014-3489-2
27. Ushirozako H, Hasegawa T, Yamato Y, et al. Impact of intrawound vancomycin powder on prevention of surgical site infection after posterior spinal surgery. *J Neurosurg Spine*. 2021;34(4):656–664. doi:10.3171/2020.8.SPINE20992
28. Tailaiti A, Shang J, Shan S, Muheremu A. Effect of Intraoperative vancomycin application in spinal surgery on the incidence of surgical site infection: a meta-analysis. *Ther Clin Risk Manag*. 2018;14:2149–2159. doi:10.2147/TCRM.S185296

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Disclosures: Gregory Grabowski reports payment/honoraria from DePuy Synthes (AO Faculty Honoraria); support for attending the AO Foundation meeting; and a leadership/fiduciary role for the Southern Orthopedic Society (board member). The remaining authors have nothing to report.

Corresponding Author: Yianni Bakaes, University of South Carolina School of Medicine Columbia, 2 Medical Park Suite 404, Columbia, SC 29203, USA; Ybakaes15@gmail.com

Published 12 October 2023

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2023 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.