

Evolving Role of Lumbar Decompression: A Narrative Review

Sagar Telang, Sahil S. Telang, Ryan Palmer, Andy Ton, William J. Karakash, Jonathan Ragheb, Siddharth Patel, Jeffrey C. Wang, Ram K. Alluri and Raymond J. Hah

Int J Spine Surg published online 24 February 2025
<https://www.ijssurgery.com/content/early/2025/02/24/8702>

This information is current as of February 24, 2025.

Email Alerts Receive free email-alerts when new articles cite this article. Sign up at:
<http://ijssurgery.com/alerts>

Evolving Role of Lumbar Decompression: A Narrative Review

SAGAR TELANG, BS¹; SAHIL S. TELANG, BS¹; RYAN PALMER, BS¹; ANDY TON, MD¹; WILLIAM J. KARAKASH, BS¹; JONATHAN RAGHEB, BS²; SIDDHARTH PATEL, BS¹; JEFFREY C. WANG, MD¹; RAM K. ALLURI, MD¹; AND RAYMOND J. HAH, MD¹

¹Department of Orthopedic Surgery, Keck School of Medicine of the University of Southern California, Los Angeles, CA, USA; ²Department of Orthopedic Surgery, Kaiser Permanente Bernard J. Tyson School of Medicine, Los Angeles, CA, USA

ABSTRACT

Traditional open lumbar decompression techniques have long been used to relieve spinal canal pressure caused by lumbar spinal stenosis. However, these procedures are associated with significant postoperative pain and prolonged recovery. Over the past few decades, there has been a shift toward minimally invasive surgical (MIS) techniques designed to minimize tissue trauma, postoperative pain, and recovery time. These advancements represent a major step forward, offering smaller incisions and direct visualization of the spinal canal. Despite the clear benefits of MIS and endoscopic techniques, they also present challenges such as a steep learning curve for surgeons and a risk of incomplete decompression. The present review examines the historical progression from open to MIS and endoscopic lumbar decompression techniques, assessing their clinical outcomes, benefits, and limitations. It highlights the ongoing need for careful application of these methods based on individual patient factors and emphasizes the importance of balancing innovative techniques with evidence-based practices to enhance patient care in spine surgery. The future of lumbar decompression will likely be shaped by further technological advancements, including navigation systems, robotic assistance, and augmented reality, which promise to improve surgical precision and outcomes.

Lumbar Spine

Keywords: endoscopic spine surgery, lumbar decompression, minimally invasive surgery, spinal stenosis, surgical techniques

INTRODUCTION

Lumbar spinal stenosis represents a substantial proportion of surgically managed spinal pathologies, particularly among the aging population.¹ Among operatively managed cases, direct lumbar decompression techniques (ie, laminectomy, laminotomy, laminoplasty, and foraminotomy) have emerged as a foundational treatment paradigm, offering less invasive approaches under select circumstances, as low-grade degenerative lumbar spondylolisthesis, relative to procedures involving fusion.^{1–4} Therefore, the capacity to minimize surgical invasiveness within lumbar decompression procedures has been a point of interest for ongoing research.

Over the past few decades, the landscape of lumbar decompression techniques has witnessed substantial evolution marked by the gradually increasing popularity of minimally invasive surgical (MIS) techniques compared with conventional, open decompression procedures. Traditionally, open decompression requires extensive dissection and retraction to access and decompress affected spinal segments.^{5–7} The advent of MIS techniques, propelled by surgical innovation

and refinements in technical expertise, has engendered approaches to minimize tissue trauma, reduce postoperative pain, and accelerate functional recovery.^{8,9} Additionally, the introduction of endoscopic spine surgery represents a further advancement, employing minimal incisions and specialized instruments for direct visualization and decompression of the spinal canal, lending a promising prospect to further refine clinical outcomes.^{10–12}

Despite the seemingly apparent benefits of MIS and endoscopic techniques, their adoption must be approached judiciously. Challenges such as a steep learning curve for surgeons, limited visualization of the surgical field, and the potential for incomplete decompression highlight the need for careful patient selection and surgeon expertise.^{13–16} Moreover, the long-term outcomes and comparative effectiveness of these techniques vs traditional approaches remain areas of active investigation.^{17,18} By critically examining the evolution of lumbar decompression techniques, from open to minimally invasive and endoscopic methods, this review aims to shed light on the progress achieved thus far and the considerations necessary for future advancements. This historical perspective underscores the importance

of balancing innovation with evidence-based practice to ensure that new techniques provide tangible benefits over established methods, ultimately enhancing patient care in the field of spine surgery. A general overview of open and minimally invasive lumbar decompression techniques is provided in Table 1.

OPEN LUMBAR DECOMPRESSION

Laminectomy

Laminectomies constitute 1 of the most common procedures used to achieve lumbar decompression, classically involving the removal of the lamina and spinous process lateral to the medial component of the facet joint at the impacted level.¹⁹ The first lumbar laminectomy was performed in 1814 by Henry Cline through an open approach, marking the foremost modern attempt at achieving spinal decompression.³³ Confronted by poor aseptic techniques and limited anesthetic capabilities, the procedure invariably failed. In 1829, however, Alban Gilpin Smith performed the first successful bilateral laminectomy to address a traumatic disc rupture.^{34,35} Following this historical milestone, open bilateral laminectomies became a foundational technique for lumbar decompression.^{35,36} In 1910, Alfred Taylor introduced a less invasive derivative, an open unilateral laminectomy, as an evolution of its bilateral counterpart.³⁷ Thereon, open unilateral approaches became the customary decompression technique practiced throughout the early and mid-1900s.³⁸

Laminoplasty

Laminoplasty has become 1 of the preferred methods for posterior decompression and is generally considered a safe and reliable procedure.³⁹ Developed by Japanese orthopedic surgeons in the 1970s and 1980s to counteract limitations with posterior laminectomy in the cervical spine, including intraoperative spinal cord injury, formation of a “laminectomy membrane,” and postoperative progression of cervical kyphosis, this technique has gained widespread popularity due to its minimally invasive approach, decreased postoperative pain, and lower risk of postoperative deformity compared with traditional laminectomy.^{40,41}

Laminoplasty has evolved through 4 primary techniques: Z-shaped (1968), open-door (1977), en-bloc (1978), and double-door or “French-Door” (1982).^{42–45} While these techniques were originally developed to treat cervical spine pathology, many of them have been adopted to treat lumbar pathology as well. Introduced in 1981, lumbar laminoplasty aimed to circumvent

the iatrogenic instability and tissue damage that often resulted from conventional laminectomies. In the late 1980s and 1990s, several studies comparing outcomes of open lumbar laminoplasty with conventional laminectomies found lower reoperation rates within 4 years for patients who underwent laminoplasty, but at the cost of increased operative time and blood loss, suggesting its benefits were mainly for younger patients with lumbar stenosis.^{20,46,47} When comparing various laminoplasty techniques, to date, no particular method of laminoplasty has yielded superior results. Especially in the lumbar spine, data comparing postoperative outcomes between the variations remain sparse.^{48,49}

In most cases, laminoplasty and laminotomy may be used interchangeably when treating posterior cord compression. A systematic review and meta-analysis performed by Sun et al suggests that laminoplasty may be preferable over laminectomy due to decreased blood loss, quicker recovery, shorter length of stay (LOS), lower rates of spinal deformity, and reduced cerebrospinal fluid leakage, without impacting operative time or tumor resection rates.⁵⁰ However, these findings must be taken with caution considering that limited high-quality data exist comparing these various techniques.

Laminotomy

Laminotomies present an alternative decompressive technique to relieve stenosis by either unilateral or bilateral removal of the upper part of the inferior lamina and the lower portion of the superior lamina.²¹ The first cited case of a laminotomy was described in the early 20th century by Hermann Oppenheim, who proposed the use of a partial laminectomy. By the 1980s, laminotomies began to be recognized as a viable alternative to laminectomies, offering the advantages of decompression of the affected area while preserving posterior structures.^{22,51,52} By preserving the supraspinous ligament, interspinous ligament, and spinous process intact, postoperative weakness associated with paraspinal denervation is minimized, and midline spinal elements vital for spinal stability are maintained.^{51,52} Unlike laminectomies, laminotomies involve reduced disruption of the surrounding biomechanics and anatomy of the spine.⁵³ The reduced invasiveness of laminotomies mitigates the risk of iatrogenic mechanical instability and has the potential to obviate the need for subsequent operative interventions when appropriately indicated.⁵⁴

Research evaluating postoperative outcomes has shown that laminotomies yield similar clinical outcomes to those of laminectomies. In a large prospective study conducted between 1985 and 1987, it was found that patients

Table 1. Summary features of lumbar decompression techniques.^{19–32}

Procedure	Summary Features
Laminectomy	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Symptomatic spinal canal stenosis that does not respond to conservative treatment ● Rapidly progressing neurological deficits or intolerable pain ● Cauda equina syndrome <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Patients with multiple medical comorbidities including depression ● Scoliosis ● Spondylolysis ● Lateral listhesis <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Open <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● High success rate with significant alleviation of preoperative symptoms ● Low rate of postoperative complications that continues to reduce with the implementation of less invasive approaches <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Potential iatrogenic disruption of posterior structures and anatomy/biomechanics of spine ● May require concomitant fusion
Laminoplasty	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Symptomatic stenosis without any signs of significant instability <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Severe osteoporosis or active infection ● Significant instability requiring fusion <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Open <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Posterior spinal structures preserved ● Reduced risk of postoperative spinal instability and deformity <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Limited literature looking at long-term outcomes of lumbar laminoplasty compared to other procedures ● Greater risk for nerve root injury intraoperatively ● Longer operative time
Laminotomy	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Spinal stenosis with neurological symptoms ● Multilevel spondylotic lateral canal stenosis ● Lateral recess stenosis secondary to disc herniation <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Severe instability necessitating fusion ● Pan-canal or central stenosis ● Severe facet joint arthritis or osteoporosis <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Open, minimally invasive, endoscopic (includes inside-out, outside-in, contralateral techniques) ● Unilateral and bilateral <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Reduced risk for iatrogenic mechanical instability and prolonged recovery when compared to laminectomy ● Superior preservation of posterior spinal structures <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Limited extent of decompression when compared to laminectomy ● May not fully alleviate central canal stenosis
Foraminotomy	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Foraminal stenosis with compression of neural structures <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Severe spinal instability (e.g., scoliosis, spondylolisthesis, kyphosis) necessitating fusion ● Significant neurological deficits requiring more extensive decompression ● Active infection or severe osteoporosis <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Open, minimally invasive, endoscopic (includes transforaminal and laser assisted endoscopic foraminotomies) <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Preserves lamina integrity ● Potential postoperative improvement of both foraminal diameter and height <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Risk of nerve injury ● Reduced access to central canal leading to potential risk for incomplete decompression and symptom recurrence

Table 1. Continued.

Procedure	Summary Features
Percutaneous Minimally Invasive Lumbar Decompression (MILD)	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Stenosis secondary to ligamentum flavum hypertrophy ≥ 2.5 mm ● Typically reserved for patients with multiple medical comorbidities making them high-risk candidates for traditional procedures <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Previous spinal surgery ● Infection at the site of potential surgery <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Percutaneous approach through a small incision ● Performed utilizing local anesthesia with the assistance of fluoroscopic guidance <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Is safe for patients with comorbidities that make it challenging to tolerate traditional spinal surgery such as laminectomy and laminotomy ● Does not require general anesthesia ● Minimally invasive, avoiding large incisions and lamina removal <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Limited decompression compared to traditional surgical method ● Limited evidence and literature available showing long-term efficacy

undergoing laminotomies experienced a 90.1% rate of symptom relief compared with 80%–90% for traditional laminectomies.²³ Further investigations have demonstrated that laminotomies yield comparable pain reduction, postoperative stability, and range of motion to laminectomies.^{24,53,54} The incidence of common procedural complications, such as neural injury, infection, durotomies, and epidural hematomas, remains similar between patients undergoing either laminectomies or laminotomies.⁵³ Laminotomies are a viable option for patients presenting with multilevel spondylotic lateral canal stenosis, who may have equal or greater levels of symptom relief through multilevel laminotomies when compared with a traditional laminectomy. However, for those patients presenting with pan-canal or central stenosis, partial decompression offered by laminotomies would not be sufficient and would necessitate a full laminectomy. The high success rate of laminotomies along with the reduced impact to midline spinal elements offer a less invasive but equally efficacious surgical approach in relieving symptoms of spinal stenosis.²³

Foraminotomy

Foraminotomies constitute a category of lumbar decompressive procedures designed to relieve compression or pressure on spinal nerves in the neural foramen.^{25,55} Distinct from laminectomies and discectomies, foraminotomies aim to preserve lamina integrity while providing precise relief from nerve compression.⁵⁶ This surgical approach is indicated for foraminal stenosis, a condition characterized by compression of a nerve root within the middle zone of the foramina, aiming to effectively address this targeted area of nerve impingement.^{57,58} Although spinal stenosis has been recognized since the early 1900s, it was not until the 1970s that a subset of patients with

lumbar nerve root compression due to superior articular facet hypertrophy were identified, leading to the adoption of foraminotomies as a viable treatment option.

MINIMALLY INVASIVE LUMBAR DECOMPRESSION

While decompression was traditionally achieved through an open approach, the field of spine surgery underwent a paradigm shift in 1994 when Foley and Smith introduced the use of tubular retractors to facilitate the treatment of lumbar pathologies.⁵⁹ The integration of tubular retractors enabled surgeons to adopt a minimally invasive approach to help alleviate spinal pathology. Tubular retractors allow surgeons to systematically expose regions they need to access through the progressive dilation of soft tissue structures without having to directly make incisions through muscle, reducing the risk of chronic postoperative back pain due to intraoperative soft tissue damage.⁶⁰

Minimally Invasive Laminotomy

The approach to laminotomies has gradually evolved, initially from an open approach, termed a classic laminotomy, to a more minimally invasive approach. The inception of unilateral and bilateral minimally invasive laminotomies,^{23,53,54,61,62} in comparison to open approaches, has demonstrated shorter LOS, greater stability, lower wound complications, less estimated blood loss, postoperative opioid use, and shorter recovery times to baseline.^{26,63} However, some studies have shown that while patients with minimally invasive laminotomies have a reduced risk of complications, there is not a significant difference in outcomes between open and MIS approaches.⁶³ While clinical outcomes are comparable between unilateral and bilateral approaches,

unilateral laminotomies have been associated with lower estimated blood loss, shorter operative times, and improved postoperative radiographic stability, the latter of which may be attributed to the intervertebral translational motion induced by bilateral procedures.⁵⁴ Unilateral laminotomies also present a marginally higher risk of postoperative complications, such as epidural hematomas, incidental durotomies, and radicular deficits.^{24,54} The increased rate of epidural hematomas may be attributed to decreased control of the contralateral epidural veins, while the increased rate of incidental durotomies has been attributed to the increased technical demand of the unilateral approach.⁵⁴

Despite these challenges, minimally invasive unilateral laminotomies for bilateral decompression (ULBD) have increasingly become the preferred choice among surgeons due to its ability to provide adequate decompression while minimizing tissue dissection and trauma, decreasing operating time, and reducing damage to spinal structures and musculature.^{53,64,65} While minimally invasive ULBDs have been associated with favorable patient outcomes, they are not without their limitations, including the need to dissect paravertebral muscles as well as poor visualization of the contralateral visual field.

Percutaneous Minimally Invasive Lumbar Decompression

Percutaneous minimally invasive lumbar decompression (MILD) was developed in 2005 by Drs. Solsberg and Schomer for patients with spinal stenosis who were unable to tolerate spinal surgery due to existing comorbidities.²⁷ MILD, which is conducted primarily with local anesthesia, offers a less invasive option compared with traditional laminotomies/laminectomies, involving the removal of portions of hypertrophied ligamentum flavum with a bone rongeur through a percutaneous port without the installation of implants. MILD is customarily preferred in patients with limited physiological resilience to more invasive procedures and is typically indicated for patients with imaging-confirmed stenosis secondary to ligamentum flavum hypertrophy ≥ 2.5 mm.²⁷ Contraindications include previous spinal surgery and infection at the site of potential surgery.²⁷ In 2016, the success of the MiDAS ENCORE randomized controlled trial (RCT) demonstrated MILD's effectiveness over epidural steroid injections for lumbar stenosis, leading to its approval for Medicare coverage beginning in 2017.^{27,28,66}

Patients undergoing MILD have shown notable postoperative improvements in both pain and functionality. Across 5 studies, MILD patients have been shown to

have a 41% improvement in visual analog scale (VAS) scores.⁶⁷⁻⁷¹ In terms of functional improvement, MILD patients in 1 study demonstrated a postoperative improvement in Oswestry Disability Index (ODI) scores by 16.5 points at 6 weeks, 16.2 at 3 months, 15.4 at 6 months, and 14.0 at 1 year.⁶⁷ MILD patients also demonstrated a low risk of adverse events intraoperatively, with the most common surgical complications being bleeding requiring blood transfusions and dural tears.⁷²

With its efficacy and safety better understood, there has been an increase in the use of MILD as an early, first-line operative treatment strategy following failed conservative treatment for patients who are poor surgical candidates. With the expanding application of MILD, future research is needed to corroborate its utility against contemporary and customary treatment strategies.²⁷ Currently, the MOTION study, an RCT involving 150 patients, is assessing the efficacy of MILD compared with conventional medical management, while another prospective study is comparing the outcomes of patients treated with MILD compared with spacers. In addition to further comparative studies, improvements in the navigation technology and optical imaging used during MILD procedures will help increase its efficacy for treating lumbar spinal stenosis.²⁷

ENDOSCOPIC LUMBAR DECOMPRESSION

In the 1980s, Parviz Kambin led the originative preliminary explorations of endoscopic approaches to lumbar spine surgery.⁷³ Endoscopic approaches have garnered substantial traction over the past decade with reduced operative times, enhanced visualization, reduced risk of paravertebral muscle injury, improved preservation of bony structures, less perioperative pain, and enhanced functional recovery.⁷⁴⁻⁷⁸ These approaches typically employ interlaminar or transforaminal techniques, utilizing uniportal, biportal, or endoscopic systems.⁷⁹ Notably, the utilization of endoscopic approaches for several procedures have demonstrated particularly promising outcomes, including endoscopic microdiscectomy, endoscopic unilateral laminotomy for bilateral decompression (ULBD), and transforaminal endoscopic lumbar foraminotomy (TELF; Table 2).

Endoscopic Microdiscectomy

In 1934, the first lumbar discectomy was performed utilizing an open approach to remove an intervertebral disc.⁹³ Consequently, in 1978, the introduction of an operating microscope marked a significant advancement, leading to the evolution of minimally invasive lumbar

Table 2. Summary features of endoscopic lumbar decompression.^{22,24,64,76-92}

Procedure	Summary Features
Endoscopic Microdiscectomy	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Lumbar disc herniation resulting in neurological deficits and radiculopathy not responding to conservative management <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Severe facet joint arthritis or active infection ● Severe central canal stenosis requiring extensive decompression than provided by microdiscectomy <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Interlaminar or Transforaminal <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Reduced soft tissue and bony trauma ● Smaller incision and improved visualization of the surgical site ● Shorter LOS in hospital and faster recovery <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Steep learning curve ● Potential for limited access to complex herniations or multi-level diseases increasing risk of incomplete decompression
Transforaminal Endoscopic Lumbar Foraminotomies (TELF)	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Foraminal stenosis with nerve root compression ● Lateral recess stenosis caused by disc herniation <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Necessity for more extensive decompression due to severe central canal stenosis ● Severe facet joint arthritis or active infection <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Transforaminal <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Improved visualization of neural structures along with smaller incisions resulting in reduced soft tissue trauma ● Enhanced preservation of structures around surgical site and improved postoperative spinal stability <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Reduced access to central canal ● Challenging to achieve full decompression in severe cases of foraminal stenosis
Endoscopic Unilateral Laminotomy for Bilateral Decompression (ULBD)	<p><u>Indications</u></p> <ul style="list-style-type: none"> ● Symptomatic disc herniation not responding to conservative management ● Lumbar spinal stenosis resulting in bilateral neural compression <p><u>Contraindications</u></p> <ul style="list-style-type: none"> ● Severe central canal stenosis requiring more invasive approaches to achieve adequate decompression ● Significant spinal instability such as vertebral fractures or severe spondylolisthesis ● Severe facet joint arthritis or active infection <p><u>Approaches</u></p> <ul style="list-style-type: none"> ● Typically interlaminar <p><u>Advantages</u></p> <ul style="list-style-type: none"> ● Preservation of contralateral structures due to unilateral approach for bilateral decompression ● Smaller incisions, reduced soft tissue trauma, quicker recovery, and shorter in hospital LOS <p><u>Disadvantages</u></p> <ul style="list-style-type: none"> ● Steep learning curve ● Limited access to central canal

Abbreviations: LOS, length of stay; TELF, transforaminal endoscopic lumbar foraminotomies; ULBD, unilateral laminotomy for bilateral decompression.

microdiscectomies, which demonstrated comparable outcomes to conventional open discectomies.⁹³ Further innovation occurred in the 1990s with the development of an endoscopic technique.^{22,94}

Data from current literature present promising outcomes for patients undergoing endoscopic discectomies compared with open and microsurgical approaches, as evidenced by several RCTs. Mayer and Brock conducted an RCT with 40 patients and showed that at 2 years post-operatively, those undergoing transforaminal endoscopic percutaneous lumbar discectomy displayed superior rates of symptom resolution (80% vs 65%) and higher rates of return to work (96% vs 72%) compared with patients who underwent microsurgical discectomy.⁹⁵ In an RCT

involving 143 patients comparing outcomes between patients undergoing transforaminal endoscopic discectomy (TED) and microdiscectomy, Gibson et al demonstrated that while both cohorts experienced significant improvements in all outcome measures, patients undergoing TED had significantly reduced hospital stay and affected side leg pain at 2 years.⁹⁶ However, TED patients had a 2-times higher risk of revision at 2 years compared with microdiscectomy patients. Ruetten et al, in an RCT with 178 patients, found similar ODI, VAS-back, and VAS-leg scores at 2 years with no significant difference in symptom recurrence between patients undergoing lumbar discectomies utilizing a full-endoscopic interlaminar technique and those undergoing surgery with a microsurgical

technique.⁹⁷ Furthermore, in a multicenter RCT involving 613 patients, Gadiradj et al reported significantly reduced VAS-back, ODI, and VAS-leg pain scores at 12 months postoperatively in patients undergoing full endoscopic discectomy compared with open discectomy for sciatica.⁸⁴

Although current literature shows promising results for endoscopic microdiscectomies compared with open and microsurgical techniques, further high-quality prospective studies are needed to elucidate long-term outcomes.

Endoscopic ULBD

While minimally invasive ULBD has shown promising outcomes, it still requires the dissection of paravertebral muscle and provides suboptimal visualization of the contralateral visual field. To overcome deficiencies, there has been an attempt to integrate the use of an endoscope for ULBDs.^{64,85–87} Endoscopic ULBDs can be subdivided into different approaches: inside-out, outside-in, and contralateral.^{80,81}

Compared with minimally invasive ULBDs, several studies have shown the benefits of endoscopic ULBD, including reduced hospital LOS and improved patient-reported outcomes such as lower ODI and VAS leg pain scores at 1 year.^{82,83,98} In a retrospective study of 93 patients, with 42 undergoing endoscopic ULBD and 51 undergoing minimally invasive ULBD, Chen et al found that patients who underwent endoscopic ULBD experienced significantly reduced VAS scores for back pain, less blood loss, shorter hospitalization, and decreased analgesic use.⁹⁸ Similarly, in another retrospective study of 60 patients comparing outcomes between patients who underwent endoscopic vs minimally invasive ULBD, Kim et al demonstrated significant postoperative improvements in ODI, VAS, and MacNab scores in both cohorts, with greater improvement in the endoscopic group.⁸³

The widespread adoption of endoscopic ULBDs as a standard practice has encountered several obstacles, including the demand for extensive training, longer operating room times, and specialized equipment.^{64,82,99} Despite these challenges, endoscopic ULBDs have shown notable advantages over conventional open and minimally invasive laminotomies, including reduced LOS and improved postoperative VAS leg pain scores and back pain disability index scores.^{80,81,100} The increasing preference for endoscopic ULBDs has spurred ongoing technical advancements, including the improvement of endoscopic systems designed to mitigate disruption of the interspinous ligament and facet

joints.^{53,100} These advancements further contribute to the procedure's evolving efficacy and safety profile.

Transforaminal Endoscopic Lumbar Foraminotomies

Similar to other procedures, endoscopic foraminotomies have gained increased popularity compared with traditional approaches to foraminotomy due to their high effectiveness and superior preservation of soft tissue and bony structures. Specifically, TELF enables exploration of the intervertebral foramen, facilitating the ablation of epidural fibrosis, resection of herniated intervertebral discs, and removal of osteophytes.^{25,101} Furthermore, TELF is recognized as a safe alternative to fusion for patients suffering concurrently from spondylolisthesis and foraminal stenosis.⁵⁵

TELF presents a less invasive solution for spinal nerve root compression in conditions such as lumbar foraminal stenosis and herniated nucleus pulposus.^{56,57} Studies have reported TELFs to have a success rate of 78%–90% with a low risk of adverse events, which is attributed to postoperative increases in foraminal height and area, thus reducing nerve compression.^{57,58} However, their effectiveness varies depending on the type of radiculopathy and location of foraminal stenosis.¹⁰² In a retrospective review of 220 patients, Lewandrowski et al found less favorable outcomes for entry zone foraminal stenosis, likely attributable to the predisposition of the entry zone to exhibit “hypertrophy of the superior articular facet.”^{29,102} Consequently, TELFs are most beneficial for middle and exit zone foraminal stenosis and contained herniated discs.

Laser-assisted endoscopic foraminotomies represent a novel technique, enhancing the precision of bone and tissue removal. Early case series show promising results, with improvements observed in both VAS and ODI scores.¹⁰³ Its highly targeted approach to decompression makes TELF a viable alternative to fusion surgery or open foraminal decompression, especially for older adults or medically compromised patients, as it does not require a repeat posterior approach and can be done under local anesthesia.⁵⁷ Nevertheless, comparative studies on endoscopic foraminotomy outcomes for lumbar decompression are limited, underscoring the need for further investigation.

Limitations of Endoscopic Approaches

Despite several notable benefits, including faster recovery times and reduced iatrogenic injury to surrounding soft tissue and bony structures, endoscopic approaches also have a few notable limitations,

including the need for specialized training and equipment. Additionally, both open and minimally invasive approaches provide distinct advantages depending on patient pathology and medical history. For instance, an endoscopic approach is preferred for laminotomies when treating patients with smaller herniations, less severe stenosis, or those with contraindications to open surgery. Conversely, open laminotomies are favored for larger lumbar herniations or more severe spinal stenosis, where extensive tissue or bone removal is necessary to achieve adequate decompression.⁸⁸

While promising outcomes associated with endoscopic techniques have resulted in advocacy for their use over minimally invasive and open approaches, the intense training and specialized equipment required for endoscopic surgery represent a limiting factor.²² The adoption of endoscopic techniques has progressed more rapidly in Asia, where a large proportion of procedures are now conducted using an endoscopic approach.²² However, there has been a delayed transition in the United States, attributed in part to increased regulations preventing trials needed to assess the efficacy of these techniques. Furthermore, it is important to consider challenges, including increased capital cost, decreased intraoperative visualization, and the steep learning curve required to effectively implement endoscopic spine surgery.²²

FUTURE OF LUMBAR DECOMPRESSION

In the coming years, the demand for lumbar decompression surgery is expected to rise. The use of endoscopic techniques is most prevalent in Asia, with over 80% of the current literature on endoscopic lumbar surgical approaches coming from the region. However, the United States and other regions have been slower to adopt these techniques due in part to increased regulations.²² The future of lumbar spine surgery appears to be moving away from traditional open approaches and toward endoscopic methods. The progression of this shift is largely attributed to technological innovations, specifically the adoption of navigation systems. Navigation systems provide surgeons with more accurate intraoperative anatomical guidance than standard 2-D fluoroscopy and enhance the accuracy of instrument placement and pedicle screw fixation. The transition from optic to electromagnetic navigation systems is also promising, reducing radiation exposure and operation time without compromising outcomes.^{84,96,97}

Robotic-assisted surgery is another area expected to grow in the coming years, potentially easing the learning curve for endoscopic surgery. Robots are capable of accurately placing endoscopes, helping with preoperative mapping and assisting with pedicle screw fixation, leading to improved patient outcomes.^{80,84-87} Several recent studies have even shown improved outcomes for patients with pedicle screw fixation carried out in surgeries with robots compared with surgeries without robots.⁸⁹⁻⁹² For laminotomies, robotic assistance during surgery has been shown to minimize human error by helping plan the laminotomy map and entry point, identifying the location of the interlaminar window, and executing precise drilling depths.¹⁰⁴

Augmented and virtual reality are anticipated to transform spine surgery even further. Virtual reality systems can improve surgical planning and training, while augmented reality offers real-time, head-mounted displays that combine navigation data with operative views to optimize surgical safety and efficiency.⁸⁹ Furthermore, augmented reality has been shown to aid in pedicle screw fixation, including helping to find entry points and ideal trajectories similar to standard navigation systems.^{89,92,105,106} There are currently 3 augmented reality devices on the market for spine surgery, including Augmedics XVision, Microsoft HoloLens, and Immersive Touch.⁹²

CONCLUSION

Lumbar decompression, driven by advancements in technology and surgical training, will continue to evolve. The integration of new technologies like navigation systems, robotic assistance, and augmented reality is expected to refine current procedures, optimize treatment, and improve patient outcomes.

REFERENCES

1. Diwan S, Sayed D, Deer TR, Salomons A, Liang K. An algorithmic approach to treating lumbar spinal stenosis: an evidenced-based approach. *Pain Med.* 2019;20(Supplement_2):S23-S31. doi:10.1093/pm/pnz133
2. Lurie J, Tomkins-Lane C. Management of lumbar spinal stenosis. *BMJ.* 2016;352:h6234. doi:10.1136/bmj.h6234
3. Katz JN, Zimmerman ZE, Mass H, Makhni MC. Diagnosis and management of lumbar spinal stenosis: a review. *JAMA.* 2022;327(17):1688-1699. doi:10.1001/jama.2022.5921
4. Genevay S, Atlas SJ. Lumbar spinal stenosis. *Best Pract Res Clin Rheumatol.* 2010;24(2):253-265. doi:10.1016/j.berh.2009.11.001
5. Nerland US, Jakola AS, Solheim O, et al. Minimally invasive decompression versus open laminectomy for central stenosis of

- the lumbar spine: pragmatic comparative effectiveness study. *BMJ*. 2015;350:h1603. doi:10.1136/bmj.h1603
6. Shih P, Wong AP, Smith TR, Lee AI, Fessler RG. Complications of open compared to minimally invasive lumbar spine decompression. *J Clin Neurosci*. 2011;18(10):1360–1364. doi:10.1016/j.jocn.2011.02.022
 7. Rahman M, Summers LE, Richter B, Mimran RI, Jacob RP. Comparison of techniques for decompressive lumbar laminectomy: the minimally invasive versus the “classic” open approach. *Minim Invasive Neurosurg*. 2008;51(2):100–105. doi:10.1055/s-2007-1022542
 8. Lawrence MM, Hayek SM. Minimally invasive lumbar decompression: a treatment for lumbar spinal stenosis. *Curr Opin Anesthesiol*. 2013;26(5):573. doi:10.1097/01.aco.0000432520.24210.54
 9. Tredway TL. Minimally invasive lumbar decompression. *Neurosurg Clin N Am*. 2006;17(4):467–476. doi:10.1016/j.nec.2006.06.003
 10. Jang JW, Lee DG, Park CK. Rationale and advantages of endoscopic spine surgery. *Int J Spine Surg*. 2021;15(suppl3):S11–S20. doi:10.14444/8160
 11. Hasan S, Härtl R, Hofstetter CP. The benefit zone of full-endoscopic spine surgery. *J Spine Surg*. 2019;5(Suppl 1):S41–S56. doi:10.21037/jss.2019.04.19
 12. Choi G, Pophale CS, Patel B, Uniyal P. Endoscopic spine surgery. *J Korean Neurosurg Soc*. 2017;60(5):485–497. doi:10.3340/jkns.2017.0203.004
 13. Lewandrowski K-U, Telfeian AE, Hellinger S, et al. Difficulties, challenges, and the learning curve of avoiding complications in lumbar endoscopic spine surgery. *Int J Spine Surg*. 2021;15(suppl 3):S21–S37. doi:10.14444/8161
 14. Sclafani JA, Kim CW. Complications associated with the initial learning curve of minimally invasive spine surgery: a systematic review. *Clin Orthop Relat Res*. 2014;472(6):1711–1717. doi:10.1007/s11999-014-3495-z
 15. Sharif S, Afsar A. Learning curve and minimally invasive spine surgery. *World Neurosurg*. 2018;119:472–478. doi:10.1016/j.wneu.2018.06.094
 16. Wang H, Huang B, Li C, et al. Learning curve for percutaneous endoscopic lumbar discectomy depending on the surgeon’s training level of minimally invasive spine surgery. *Clin Neurol Neurosurg*. 2013;115(10):1987–1991. doi:10.1016/j.clineuro.2013.06.008
 17. Sen RD, White-Dzuro G, Ruzevick J, et al. Intra- and peri-operative complications associated with endoscopic spine surgery: a multi-institutional study. *World Neurosurg*. 2018;120:e1054–e1060. doi:10.1016/j.wneu.2018.09.009
 18. Kim HS, Wu PH, Jang IT. Current and future of endoscopic spine surgery: what are the common procedures we have now and what lies ahead? *World Neurosurg*. 2020;140:642–653. doi:10.1016/j.wneu.2020.03.111
 19. Estefan M, Munakomi S, Camino Willhuber GO. Laminectomy. In: *StatPearls*. StatPearls Publishing; 2024. <http://www.ncbi.nlm.nih.gov/books/NBK542274/>. Accessed March 26, 2024.
 20. Tsuji H, Itoh T, Sekido H, et al. Expansive laminoplasty for lumbar spinal stenosis. *Int Orthop*. 1990;14(3):309–314. doi:10.1007/BF00178765
 21. Ruggeri A, Pichierri A, Marotta N, Tarantino R, Delfini R. Laminotomy in adults: technique and results. *Eur Spine J*. 2012;21(2):364–372. doi:10.1007/s00586-011-1826-2
 22. Mayer HM. A history of endoscopic lumbar spine surgery: what have we learnt? *Biomed Res Int*. 2019;2019:4583943. doi:10.1155/2019/4583943
 23. Aryanpur J, Ducker T. Multilevel lumbar laminotomies: an alternative to laminectomy in the treatment of lumbar stenosis. *Neurosurgery*. 1990;26(3):429–432.
 24. Ho YH, Tu YK, Hsiao CK, Chang CH. Outcomes after minimally invasive lumbar decompression: a biomechanical comparison of unilateral and bilateral laminotomies. *BMC Musculoskeletal Disord*. 2015;16:208. doi:10.1186/s12891-015-0659-2
 25. Ahn Y, Oh HK, Kim H, Lee SH, Lee HN. Percutaneous endoscopic lumbar foraminotomy: an advanced surgical technique and clinical outcomes. *Neurosurgery*. 2014;75(2):124–133. doi:10.1227/NEU.0000000000000361
 26. Tan LA, Kasliwal MK, Fessler RG. Minimally invasive versus open laminotomy. *Spine J*. 2014;14(6):1081–1082. doi:10.1016/j.spinee.2013.12.030
 27. Jain S, Deer T, Sayed D, et al. Minimally invasive lumbar decompression: a review of indications, techniques, efficacy and safety. *Pain Manag*. 2020;10(5):331–348. doi:10.2217/pmt-2020-0037
 28. Staats PS, Benyamin RM, Investigators ME. MiDAS encore: randomized controlled clinical trial report of 6-month results. *Pain Physician*. 2016;19(2):25–38.
 29. Lee CK, Rauschnig W, Glenn W. Lateral lumbar spinal canal stenosis: classification, pathologic anatomy and surgical decompression. *Spine (Phila Pa 1976)*. 1988;13(3):313–320. doi:10.1097/00007632-198803000-00015
 30. Chen Z, Zhang L, Dong J, et al. Percutaneous transforaminal endoscopic discectomy versus microendoscopic discectomy for lumbar disc herniation: two-year results of a randomized controlled trial. *Spine (Phila Pa 1976)*. 2020;45(8):493–503. doi:10.1097/BRS.0000000000003314
 31. Chen Z, Zhang L, Dong J, et al. Percutaneous transforaminal endoscopic discectomy versus microendoscopic discectomy for lumbar disc herniation: 5-year long-term results of a randomized controlled trial. *Spine (Phila Pa 1986)*. 2023;48(2):79–88. doi:10.1097/BRS.0000000000004468
 32. Sivakanthan S, Hasan S, Hofstetter C. Full-endoscopic lumbar discectomy. *Neurosurg Clin N Am*. 2020;31(1):1–7. doi:10.1016/j.nec.2019.08.016
 33. Oliver M, Zarb G, Silver J, Moore M, Salisbury V. Spinal cord injury in context. In: Oliver M, Zarb G, Silver J, Moore M, Salisbury V, eds. *Walking into Darkness: The Experience of Spinal Cord Injury*. UK: Macmillan Education UK; 1988:1–6. doi:10.1007/978-1-349-19451-3_1
 34. Emch TM, Modic MT. Imaging of lumbar degenerative disk disease: history and current state. *Skeletal Radiol*. 2011;40(9):1175–1189. doi:10.1007/s00256-011-1163-x
 35. Patchell RA, Tibbs PA, Young AB, Clark DB, Alban G. Smith and the beginnings of spinal surgery. *Neurology (Echronicon)*. 1987;37(10):1683–1683. doi:10.1212/WNL.37.10.1683
 36. Ahmed SI, Javed G, Bareeqa SB, et al. Comparison of decompression alone versus decompression with fusion for stenotic lumbar spine: a systematic review and meta-analysis. *Cureus*. 2018;10(8):e3135. doi:10.7759/cureus.3135
 37. Taylor AS. Unilateral laminectomy. *Ann Surg*. 1910;51(4):529–533. doi:10.1097/00000658-191004000-00010
 38. Oppenheim H, Krause F. Ueber einklemmung bzw. strangulation der cauda equina. *Dtsch Med Wochenschr*. 1909;35(16):697–700. doi:10.1055/s-0029-1201407

39. Hardman J, Graf O, Kouloumberis PE, Gao WH, Chan M, Roitberg BZ. Clinical and functional outcomes of laminoplasty and laminectomy. *Neurol Res.* 2010;32(4):416–420. doi:10.1179/174313209X459084
40. Ito M, Nagahama K. Laminoplasty for cervical myelopathy. *Global Spine J.* 2012;2(3):187–193. doi:10.1055/s-0032-1315456
41. Millward CP, Bhagawati D, Chan HW, Bestwick J, Brecknell JE. Retrospective observational comparative study of hemilaminectomy versus laminectomy for intraspinal tumour resection; shorter stays, lower analgesic usage and less kyphotic deformity. *Br J Neurosurg.* 2015;29(3):390–395. doi:10.3109/02688697.2014.1003026
42. Oyama M, Hattori S, Moriwaki N. A new method of cervical laminectomy. *Chubu Nippon Seikeisaigaijeka Gakkai Zasshi.* 1973;16:792–794.
43. Miyazaki K, Kirita Y. Extensive simultaneous multisegment laminectomy for myelopathy due to the ossification of the posterior longitudinal ligament in the cervical region. *Spine (Phila Pa 1976).* 1986;11(6):531–542. doi:10.1097/00007632-198607000-00005
44. Hirabayashi K, Watanabe K, Wakano K, Suzuki N, Satomi K, Ishii Y. Expansive open-door laminoplasty for cervical spinal stenotic myelopathy. *Spine (Phila Pa 1976).* 1983;8(7):693–699. doi:10.1097/00007632-198310000-00003
45. Kurokawa T, Tsyuyama M, Tanaka K, Kurokawa T, Tsyuyama N, Tanaka H. Enlargement of spinal canal by the sagittal splitting of the spinous process – ScienceOpen. <https://www.scienceopen.com/document?vid=d1a74f16-99a3-42bf-a1af-4e9c6b053c2d>. Accessed December 27, 2023.
46. Matsui H, Kanamori M, Ishihara H, Hirano N, Tsuji H. Expansive lumbar laminoplasty for degenerative spinal stenosis in patients below 70 years of age. *Eur Spine J.* 1997;6(3):191–196. doi:10.1007/BF01301435
47. Kawaguchi Y, Kanamori M, Ishihara H, et al. Clinical and radiographic results of expansive lumbar laminoplasty in patients with spinal stenosis. *The Journal of Bone and Joint Surgery-American Volume.* 2004;86(8):1698–1703. doi:10.2106/00004623-200408000-00013
48. Mitsunaga LK, Klineberg EO, Gupta MC. Laminoplasty techniques for the treatment of multilevel cervical stenosis. *Adv Orthop.* 2012;2012:307916. doi:10.1155/2012/307916
49. Byvaltsev V, Polkin R, Kalinin A, et al. Laminoplasty versus laminectomy in the treatment of primary spinal cord tumors in adult patients: a systematic review and meta-analysis of observational studies. *Asian Spine J.* 2023;17(3):595–609. doi:10.31616/asj.2022.0184
50. Sun S, Li Y, Wang X, et al. Safety and efficacy of laminoplasty versus laminectomy in the treatment of spinal cord tumors: a systematic review and meta-analysis. *World Neurosurg.* 2019;125:136–145. doi:10.1016/j.wneu.2018.12.033
51. Overvest GM, Jacobs W, Vleggeert-Lankamp C, et al. Effectiveness of posterior decompression techniques compared with conventional laminectomy for lumbar stenosis. *Cochrane Database of Systematic Reviews.* 2015;2015(3). doi:10.1002/14651858.CD010036.pub2
52. Moisi M, Fisahn C, Tkachenko L, et al. Unilateral laminotomy with bilateral spinal canal decompression for lumbar stenosis: a technical note. *Cureus.* 2016;8(5):e623. doi:10.7759/cureus.623
53. Thomé C, Zevgaridis D, Leheha O, et al. Outcome after less-invasive decompression of lumbar spinal stenosis: a randomized comparison of unilateral laminotomy, bilateral laminotomy, and laminectomy. *J Neurosurg Spine.* 2005;3(2):129–141. doi:10.3171/spi.2005.3.2.0129
54. Hong S-W, Choi KY, Ahn Y, et al. A comparison of unilateral and bilateral laminotomies for decompression of L4-L5 spinal stenosis. *Spine (Phila Pa 1976).* 2011;36(3):E172–E178. doi:10.1097/BRS.0b013e3181db998c
55. Ahn Y, Park HB, Yoo BR, Jeong TS. Endoscopic lumbar foraminotomy for foraminal stenosis in stable spondylolisthesis. *Front Surg.* 2022;9:1042184. doi:10.3389/fsurg.2022.1042184
56. Kunogi J, Hasue M. Diagnosis and operative treatment of intraforaminal and extraforaminal nerve root compression. *Spine (Phila Pa 1976).* 1991;16(11):1312–1320. doi:10.1097/00007632-199111000-00012
57. Giordan E, Billeci D, Del Verme J, Varrassi G, Coluzzi F. Endoscopic transforaminal lumbar foraminotomy: a systematic review and meta-analysis. *Pain Ther.* 2021;10(2):1481–1495. doi:10.1007/s40122-021-00309-1
58. Evins AI, Banu MA, Njoku I, et al. Endoscopic lumbar foraminotomy. *J Clin Neurosci.* 2015;22(4):730–734. doi:10.1016/j.jocn.2014.10.025
59. Prabhu MC, Jacob KC, Patel MR, Pawlowski H, Vanjani NN, Singh K. History and evolution of the minimally invasive transforaminal lumbar interbody fusion. *Neurospine.* 2022;19(3):479–491. doi:10.14245/ns.2244122.061
60. Kim YB, Hyun SJ. Clinical applications of the tubular retractor on spinal disorders. *J Korean Neurosurg Soc.* 2007;42(4):245–250. doi:10.3340/jkns.2007.42.4.245
61. Young S, Veerapen R, O’Laioire SA. Relief of lumbar canal stenosis using multilevel subarticular fenestrations as an alternative to wide laminectomy: preliminary report. *Neurosurgery.* 1988;23(5):628–633. doi:10.1227/00006123-198811000-00014
62. McCulloch J. *Microsurgical Spinal Laminotomies in the Adult Spine: Principles and Practice.* J.W. Frymoyer (ed) Raven Press.
63. Ang C-L, Phak-Boon Tow B, Fook S, et al. Minimally invasive compared with open lumbar laminotomy: no functional benefits at 6 or 24 months after surgery. *Spine J.* 2015;15(8):1705–1712. doi:10.1016/j.spinee.2013.07.461
64. McGrath LB, White-Dzuro GA, Hofstetter CP. Comparison of clinical outcomes following minimally invasive or lumbar endoscopic unilateral laminotomy for bilateral decompression. *J Neurosurg Spine.* 2019;30(4):491–499. doi:10.3171/2018.9.SPINE18689
65. Palmer S, Turner R, Palmer R. Bilateral decompression of lumbar spinal stenosis involving a unilateral approach with microscope and tubular retractor system. *J Neurosurg.* 2002;97(2 Suppl):213–217. doi:10.3171/spi.2002.97.2.0213
66. Chen H, Kelling J. Mild procedure for lumbar decompression: a review. *Pain Pract.* 2013;13(2):146–153. doi:10.1111/j.1533-2500.2012.00574.x
67. Kreiner DS, MacVicar J, Duszynski B, Nampiaparampil DE. The mild® procedure: a systematic review of the current literature: the mild® procedure: a systematic review. *Pain Med.* 2014;15(2):196–205. doi:10.1111/pme.12305
68. Brown LL. A double-blind, randomized, prospective study of epidural steroid injection vs. the mild® procedure in patients with symptomatic lumbar spinal stenosis. *Pain Pract.* 2012;12(5):333–341. doi:10.1111/j.1533-2500.2011.00518.x
69. Chopko B, Caraway DL. MiDAS I (mild decompression alternative to open surgery): a preliminary report of a prospective, multi-center clinical study. *Pain Physician.* 2010;13(4):369–378.

70. Lingreen R, Grider JS. Retrospective review of patient self-reported improvement and post-procedure findings for mild (minimally invasive lumbar decompression). *Pain Physician*. 2010;13(6):555–560.
71. Durkin B, Romeiser J, Shroyer ALW, et al. Report from a quality assurance program on patients undergoing the MILD procedure. *Pain Med*. 2013;14(5):650–656. doi:10.1111/pme.12079
72. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med*. 2008;358(8):794–810. doi:10.1056/NEJMoa0707136
73. Samartzis D, Shen FH, Perez-Cruet MJ, Anderson DG. Minimally invasive spine surgery: a historical perspective. *Orthop Clin North Am*. 2007;38(3):305–326. doi:10.1016/j.ocl.2007.04.006
74. Mobbs R, Phan K. Minimally invasive unilateral laminectomy for bilateral decompression. *JBSJ Essent Surg Tech*. 2017;7(1):e9. doi:10.2106/JBJS.ST.16.00072
75. Park S-M, Park J, Jang HS, et al. Biportal endoscopic versus microscopic lumbar decompressive laminectomy in patients with spinal stenosis: a randomized controlled trial. *Spine J*. 2020;20(2):156–165. doi:10.1016/j.spinee.2019.09.015
76. Mobbs RJ, Li J, Sivabalan P, Raley D, Rao PJ. Outcomes after decompressive laminectomy for lumbar spinal stenosis: comparison between minimally invasive unilateral laminectomy for bilateral decompression and open laminectomy: clinical article. *J Neurosurg Spine*. 2014;21(2):179–186. doi:10.3171/2014.4.SP.INE13420
77. Uehara M, Takahashi J, Hashidate H, et al. Comparison of spinous process-splitting laminectomy versus conventional laminectomy for lumbar spinal stenosis. *Asian Spine J*. 2014;8(6):768–776. doi:10.4184/asj.2014.8.6.768
78. Tang S, Mok TN, He Q, et al. Comparison of clinical and radiological outcomes of full-endoscopic versus microscopic lumbar decompression laminectomy for the treatment of lumbar spinal stenosis: a systematic review and meta-analysis. *Ann Palliat Med*. 2021;10(10):10130–10146. doi:10.21037/apm-21-198
79. Butler AJ, Alam M, Wiley K, Ghasem A, Rush Iii AJ, Wang JC. Endoscopic lumbar surgery: the state of the art in 2019. *Neurospine*. 2019;16(1):15–23. doi:10.14245/ns.1938040.020
80. Wu PH, Kim HS, Jang IT. How i do it? Uniportal full endoscopic contralateral approach for lumbar foraminal stenosis with double crush syndrome. *Acta Neurochir (Wien)*. 2020;162(2):305–310. doi:10.1007/s00701-019-04157-z
81. Kim HS, Wu PH, Jang IT. Lumbar endoscopic unilateral laminotomy for bilateral decompression outside-in approach: a proctorship guideline with 12 steps of effectiveness and safety. *Neurospine*. 2020;17(Suppl 1):S99–S109. doi:10.14245/ns.2040078.039
82. Heo DH, Lee N, Park CW, Kim HS, Chung HJ. Endoscopic unilateral laminotomy with bilateral discectomy using biportal endoscopic approach: technical report and preliminary clinical results. *World Neurosurg*. 2020;137:31–37. doi:10.1016/j.wneu.2020.01.190
83. Kim HS, Choi SH, Shim DM, Lee IS, Oh YK, Woo YH. Advantages of new endoscopic unilateral laminectomy for bilateral decompression (ULBD) over conventional microscopic ULBD. *Clin Orthop Surg*. 2020;12(3):330–336. doi:10.4055/cios19136
84. Gadjaradj PS, Rubinstein SM, Peul WC, et al. Full endoscopic versus open discectomy for sciatica: randomised controlled non-inferiority trial. *BMJ*. 2022;376:e065846. doi:10.1136/bmj-2021-065846
85. Komp M, Hahn P, Oezdemir S, et al. Bilateral spinal decompression of lumbar central stenosis with the full-endoscopic interlaminar versus microsurgical laminotomy technique: a prospective, randomized, controlled study. *Pain Physician*. 2015;18(1):61–70.
86. Hasan S, McGrath LB, Sen RD, Barber JK, Hofstetter CP. Comparison of full-endoscopic and minimally invasive decompression for lumbar spinal stenosis in the setting of degenerative scoliosis and spondylolisthesis. *Neurosurg Focus*. 2019;46(5):E16. doi:10.3171/2019.2.FOCUS195
87. Lee CW, Yoon KJ, Jun JH. Percutaneous endoscopic laminotomy with flavectomy by uniportal, unilateral approach for the lumbar canal or lateral recess stenosis. *World Neurosurg*. 2018;113:e129–e137. doi:10.1016/j.wneu.2018.01.195
88. Müller SJ, Burkhardt BW, Oertel JM. Management of dural tears in endoscopic lumbar spinal surgery: a review of the literature. *World Neurosurg*. 2018;119:494–499. doi:10.1016/j.wneu.2018.05.251
89. Kwon H, Park JY. The role and future of endoscopic spine surgery: a narrative review. *Neurospine*. 2023;20(1):43–55. doi:10.14245/ns.2346236.118
90. Gao S, Wei J, Li W, et al. Accuracy of robot-assisted percutaneous pedicle screw placement under regional anesthesia: a retrospective cohort study. *Pain Res Manag*. 2021;2021:6894001. doi:10.1155/2021/6894001
91. D'Souza M, Gendreau J, Feng A, Kim LH, Ho AL, Veeravagu A. Robotic-assisted spine surgery: history, efficacy, cost, and future trends. *Robot Surg Res Rev*. 2019;6:9–23. doi:10.2147/RSRR.S190720
92. Pierzchajlo N, Stevenson TC, Huynh H, et al. Augmented reality in minimally invasive spinal surgery: a narrative review of available technology. *World Neurosurg*. 2023;176:35–42. doi:10.1016/j.wneu.2023.04.030
93. Rasouli MR, Rahimi-Movaghar V, Shokraneh F, Moradi-Lakeh M, Chou R. Minimally invasive discectomy versus microdiscectomy/open discectomy for symptomatic lumbar disc herniation. *Cochrane Database Syst Rev*. 2014;2014(9):CD010328. doi:10.1002/14651858.CD010328.pub2
94. Perez-Cruet MJ, Foley KT, Isaacs RE, et al. Microendoscopic lumbar discectomy: technical note. *Neurosurgery*. 2002;51(5 Suppl):S129–36.
95. Mayer HM, Brock M. Percutaneous endoscopic discectomy: surgical technique and preliminary results compared to microsurgical discectomy. *J Neurosurg*. 1993;78(2):216–225. doi:10.3171/jns.1993.78.2.0216
96. Gibson JNA, Subramanian AS, Scott CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. *Eur Spine J*. 2017;26(3):847–856. doi:10.1007/s00586-016-4885-6
97. Ruetten S, Komp M, Merk H, Godolias G. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. *Spine (Phila Pa 1976)*. 2008;33(9):931–939. doi:10.1097/BRS.0b013e31816c8af7
98. Chen KT, Choi KC, Shim HK, Lee DC, Kim JS. Full-endoscopic versus microscopic unilateral laminotomy for bilateral decompression of lumbar spinal stenosis at L4-L5: comparative study. *Int Orthop*. 2022;46(12):2887–2895. doi:10.1007/s00264-022-05549-0
99. Yang JC, Kim SG, Kim TW, Park KH. Analysis of factors contributing to postoperative spinal instability after lumbar decompression for spinal stenosis. *Korean J Spine*. 2013;10(3):149–154. doi:10.14245/kjs.2013.10.3.149

100. Guiot BH, Khoo LT, Fessler RG. A minimally invasive technique for decompression of the lumbar spine. *Spine (Phila Pa 1976)*. 2002;27(4):432–438. doi:10.1097/00007632-200202150-00021

101. Knight M, Goswami A. Management of isthmic spondylolisthesis with posterolateral endoscopic foraminal decompression. *Spine (Phila Pa 1976)*. 2003;28(6):573–581. doi:10.1097/01.BRS.0000050400.16499.ED

102. Lewandrowski KU. “Outside-in” technique, clinical results, and indications with transforaminal lumbar endoscopic surgery: a retrospective study on 220 patients on applied radiographic classification of foraminal spinal stenosis. *Int J Spine Surg*. 2014;8:26. doi:10.14444/1026

103. Ahn Y, Keum HJ, Shin SH, Choi JJ. Laser-assisted endoscopic lumbar foraminotomy for failed back surgery syndrome in elderly patients. *Lasers Med Sci*. 2020;35(1):121–129. doi:10.1007/s10103-019-02803-7

104. Li Y, Wang MY. Robotic-assisted endoscopic laminotomy: 2-dimensional operative video. *Operative Surg*. 2021;20(5):E361–E361. doi:10.1093/ons/opaa441

105. Molina CA, Theodore N, Ahmed AK, et al. Augmented reality-assisted pedicle screw insertion: a cadaveric proof-of-concept study. *J Neurosurg Spine*. 2019;31(1):139–146. doi:10.3171/2018.12.SPINE181142

106. Ghaednia H, Fourman MS, Lans A, et al. Augmented and virtual reality in spine surgery, current applications and future potentials. *Spine J*. 2021;21(10):1617–1625. doi:10.1016/j.spinee.2021.03.018

Funding: The authors received no financial support for the research, authorship, and/or publication of this article.

Declaration of Conflicting Interests: The authors report no conflicts of interest in this work.

Disclosures: Sagar Telang, Sahil S. Telang, Ryan Palmer, Andy Ton, William J Karakash, Jonathan Ragheb, and Siddharth Patel have nothing to disclose. Jeffrey C. Wang has received intellectual property royalties from Zimmer Biomet, NovApproach, SeaSpine, and DePuy Synthes, and stock options from Bone Biologics, Electrocore, PearlDiver, and Surgitech. Raymond J. Hah has received grant funding from SI bone, consulting fees from NuVasive, and support from the North American Spine Society to attend meetings. Ram K. Alluri has received grant funding from NIH, consulting fees and stock options from HIA Technologies, and payment from Eccential Robotics for lectures and presentations.

Corresponding Author: William J. Karakash, Department of Orthopaedic Surgery, Keck School of Medicine of USC, 1500 San Pablo St, Los Angeles, CA 90033, USA; wkarakas@usc.edu

This manuscript is generously published free of charge by ISASS, the International Society for the Advancement of Spine Surgery. Copyright © 2025 ISASS. To see more or order reprints or permissions, see <http://ijssurgery.com>.